arXiv:1207.5347v1 [nlin.SI] 23 Jul 2012

On the τ-functions of the Degasperis-Procesi equation

Bao-Feng Feng, Ken-ichi Maruno, and Yasuhiro Ohta

Abstract. The reciprocal link between the Degasperis-Procesi (DP) equation and the pseudo 3-reduction of the C_∞ two-dimensional Toda system is used to construct the N-soliton solution of the DP equation. The N-soliton solution of the DP equation is presented in the form of pfaffian through a hodograph (reciprocal) transformation. The bilinear equations, the identities between determinants and pfaffians, and the τ-function of the DP equation are obtained from the pseudo 3-reduction of the C_∞ two-dimensional Toda system.

1. Introduction

In this article, we investigate the N-soliton solution of the Degasperis-Procesi (DP) equation [**1**]

$$u_t + 3\kappa^3 u_x - u_{txx} + 4uu_x = 3u_x u_{xx} + uu_{xxx}\,, \tag{1.1}$$

which has received much attention in recent years. Since its discovery in 1999, many interesting mathematical properties of the DP equation have been found [**2, 3, 4, 5, 6, 7, 8**]. For example, the DP equation has the N-soliton solution if $\kappa \neq 0$ [**6, 7**].

In this contribution, we show the reciprocal link between the DP equation and the pseudo 3-reduction of the C_∞ two-dimensional Toda system and investigate their τ-functions. Using this reciprocal link, we construct the N-soliton solution of the DP equation in the form of pfaffian. The bilinear equations, the identities between determinants and pfaffians, and the τ-function of the DP equation are easily obtained from the pseudo 3-reduction of the C_∞ two-dimensional Toda system.

2. The DP equation and the pseudo 3-reduction of the C_∞ 2D-Toda system

The two-dimensional Toda (2D-Toda) system of A_∞-type, which is also called the Toda field equation or the two-dimensional Toda lattice, is given as follows [**9, 10, 11, 12**]:

$$\frac{\partial^2 \theta_n}{\partial x_1 \partial x_{-1}} = -\sum_{m\in\mathbb{Z}} a_{n,m} e^{-\theta_m}\,, \quad n \in \mathbb{Z} \tag{2.1}$$

1991 *Mathematics Subject Classification.* Primary 35C08, 37K40 ; Secondary 35Q51, 37K10.

Key words and phrases. The Degasperis-Procesi equation, pfaffian, the C_∞ two-dimensional Toda system.

where the matrix $A = (a_{n,m})$ is the transpose of the Cartan matrix for the infinite dimensional Lie algebra A_∞, which is the infinite tridiagonal matrix

$$A = \begin{bmatrix} \ddots & \ddots & \ddots & & & & & \\ & -1 & 2 & -1 & & & & \\ & & -1 & 2 & -1 & & & \\ & & & \ddots & \ddots & \ddots & & \\ & & & & -1 & 2 & -1 & \\ & & & & & \ddots & \ddots & \ddots \end{bmatrix} . \tag{2.2}$$

The A_∞ 2D-Toda system (2.1) with (2.2) may be written as

$$\frac{\partial^2 \theta_n}{\partial x_1 \partial x_{-1}} = e^{-\theta_{n-1}} - 2e^{-\theta_n} + e^{-\theta_{n+1}} . \tag{2.3}$$

The A_∞ 2D-Toda system (2.1) with (2.2) is transformed into the bilinear equation

$$-\left(\frac{1}{2} D_{x_1} D_{x_{-1}} - 1\right) \tau_n \cdot \tau_n = \tau_{n-1}\tau_{n+1} , \tag{2.4}$$

through the dependent variable transformation

$$\theta_n = -\ln \frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2} . \tag{2.5}$$

Here D_x is the Hirota D-operator which is defined as

$$D_x^n a(x) \cdot b(x) = (\partial_x - \partial_{x'})^n \, a(x) b(x')|_{x=x'} . \tag{2.6}$$

Lemma 2.1 (Ueno-Takasaki[**13**], Babich-Matveev-Sall[**14**], Hirota[**15**],Nimmo-Willox[**16**]). *The bilinear equation (2.4) and other bilinear equations of the members of the 2D-Toda lattice hierarchy have the following Gram-type determinant solution:*

$$\tau_n = \det \left(\psi_{i,j}^{(n)} \right)_{1 \le i,j \le M} ,$$

where

$$\psi_{i,j}^{(n)} = c_{i,j} + (-1)^n \int_{-\infty}^{x_1} \varphi_i^{(n)} \hat{\varphi}_j^{(-n)} dx_1 .$$

Here $c_{i,j}$ *are constants,* $\varphi_{i,j}^{(n)}$ *and* $\hat{\varphi}_{i,j}^{(n)}$ *must satisfy* $\frac{\partial \varphi_i^{(n)}}{\partial x_k} = \varphi_i^{(n+k)}$ *and* $\frac{\partial \hat{\varphi}_i^{(n)}}{\partial x_k} = (-1)^{k-1} \hat{\varphi}_i^{(n+k)}$ *for* $k = \pm 1, \pm 2, \pm 3, \cdots$.

For example, the following linear independent set of functions $\{\varphi_i^{(n)}, \hat{\varphi}_j^{(n)}\}$ *for* $i, j = 1, 2, \cdots, M$ *gives* M*-soliton solution of the* A_∞ *2D-Toda system:*

$$\varphi_i^{(n)} = p_i^n e^{\xi_i} , \quad \hat{\varphi}_j^{(n)} = q_j^n e^{\eta_j} ,$$

where $\xi_i = p_i x_1 + \frac{1}{p_i} x_{-1} + p_i^2 x_2 + \frac{1}{p_i^2} x_{-2} + p_i^3 x_3 + \frac{1}{p_i^3} x_{-3} \cdots + \xi_{i0}$ *and* $\eta_i = q_i x_1 + \frac{1}{q_i} x_{-1} - q_i^2 x_2 - \frac{1}{q_i^2} x_{-2} + q_i^3 x_3 + \frac{1}{q_i^3} x_{-3} \cdots + \eta_{i0}$.

Proof. See [**15**]. □

We impose the C_∞-reduction $\theta_n = \theta_{-n}$ $(n \ge 0)$ to the A_∞ 2D-Toda system, *i.e.*, fold the infinite sequence $\{\dots, \theta_{-2}, \theta_{-1}, \theta_0, \theta_1, \theta_2, \dots\}$ in θ_0 [**17, 13, 16, 18, 19, 20**]. Then we have $\theta_{-1} = \theta_1$, $\theta_{-2} = \theta_2$, $\theta_{-3} = \theta_3$, $\dots$.

For $n=0$,

$$\begin{aligned}\frac{\partial^2\theta_0}{\partial x_1\partial x_{-1}} &= e^{-\theta_{-1}}-2e^{-\theta_0}+e^{-\theta_1}\\ &= -2e^{-\theta_0}+2e^{-\theta_1}\,.\end{aligned}$$

For $n=1$,

$$\frac{\partial^2\theta_1}{\partial x_1\partial x_{-1}} = e^{-\theta_0}-2e^{-\theta_1}+e^{-\theta_2}\,.$$

Thus we obtain the C_∞ 2D-Toda system [**9, 12**]:

$$\frac{\partial^2\theta_n}{\partial x_1\partial x_{-1}} = -\sum_{m\in\mathbf{Z}_{\geq 0}} a_{n+1\,,\,m+1}e^{-\theta_m}\,, \quad n\in\mathbf{Z}_{\geq 0}\,, \tag{2.7}$$

where the matrix $A=(a_{n,m})$ is the transpose of the Cartan matrix for the infinite dimensional Lie algebra C_∞, which is the semi-infinite tridiagonal matrix

$$A=\begin{bmatrix} 2 & -2 & & & & & \\ -1 & 2 & -1 & & & & \\ & -1 & 2 & -1 & & & \\ & & \ddots & \ddots & \ddots & & \\ & & & -1 & 2 & -1 & \\ & & & & \ddots & \ddots & \ddots \end{bmatrix}. \tag{2.8}$$

The C_∞ 2D-Toda system (2.7) with (2.8) may be written as

$$\frac{\partial^2\theta_0}{\partial x_1\partial x_{-1}} = -2e^{-\theta_0}+2e^{-\theta_1}\,, \tag{2.9}$$

$$\frac{\partial^2\theta_n}{\partial x_1\partial x_{-1}} = e^{-\theta_{n-1}}-2e^{-\theta_n}+e^{-\theta_{n+1}}\,, \qquad n\geq 1\,. \tag{2.10}$$

The C_∞ 2D-Toda system (2.7) with (2.8) is transformed into the bilinear equations

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_0\cdot\tau_0=\tau_1^2\,, \tag{2.11}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_n\cdot\tau_n=\tau_{n-1}\tau_{n+1}\,, \quad \text{for} \quad n\geq 1\,, \tag{2.12}$$

through the dependent variable transformation

$$\theta_0=-\ln\frac{\tau_1^2}{\tau_0^2}\,, \qquad \text{and} \qquad \theta_n=-\ln\frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2} \quad \text{for} \quad n\geq 1.$$

Lemma 2.2. *The bilinear equations of the C_∞ 2D-Toda system (2.11) and (2.12) have the N-soliton solution which is expressed as*

$$\tau_n=\det\left(\psi_{i,j}^{(n)}\right)_{1\leq i,j\leq 2N}\,,$$

where

$$\begin{aligned}\psi_{i,j}^{(n)} &= c_{i,j}+(-1)^n\int_{-\infty}^{x_1}\varphi_i^{(n)}\varphi_j^{(-n)}dx_1\,,\\ \varphi_i^{(n)} &= p_i^n e^{\xi_i}\,, \qquad \xi_i=p_ix_1+\frac{1}{p_i}x_{-1}+p_i^3x_3+\frac{1}{p_i^3}x_{-3}+\cdots+\xi_{i0}\,,\end{aligned}$$

and $c_{i,j}=c_{j,i}$.

Proof. Imposing the C_∞ reduction $\tau_n=\tau_{-n}$, *i.e.*, folding the sequence of the τ-functions $\{\dots,\tau_{-2},\tau_{-1},\tau_0,\tau_1,\tau_2,\dots\}$ in τ_0, we have $\tau_{-1}=\tau_1$, $\tau_{-2}=\tau_2$, $\tau_{-3}=\tau_3$, ... [**17, 13, 16, 18, 19, 20**]. Thus we obtain the bilinear equations (2.11) and (2.12) from the 2D-Toda bilinear equation (2.4).

To impose the C_∞ reduction to the Gram-type determinant solution of the A_∞ 2D-Toda system, we impose the constraint $\hat{\varphi}_j^{(n)}=\varphi_j^{(n)}$, $c_{i,j}=c_{j,i}$, $M=2N$ and $x_{2k}\equiv 0$ for every interger k. With this constraint, each element of the Gram-type determinant has the following property:

$$\begin{aligned}\psi_{i,j}^{(n)}&=c_{i,j}+(-1)^n\int_{-\infty}^{x_1}\varphi_i^{(n)}\varphi_j^{(-n)}dx_1\\&=c_{j,i}+(-1)^{-n}\int_{-\infty}^{x_1}\varphi_j^{(-n)}\varphi_i^{(n)}dx_1\\&=\psi_{j,i}^{(-n)}\,.\end{aligned}$$

Then the τ-function satisfies $\tau_n=\tau_{-n}$. Therefore the N-soliton solution of the C_∞ 2D-Toda system is expressed by the above Gram-type determinant. □

Lemma 2.3. *Consider the following τ-function:*

$$\tau_n=\det\left(\psi_{i,j}^{(n)}\right)_{1\le i,j\le 2N}\,,$$

where

$$\begin{aligned}&\psi_{i,j}^{(n)}=c_{i,j}+(-1)^n\int_{-\infty}^{x_1}\varphi_i^{(n)}\varphi_j^{(-n)}dx_1\,,\\&\varphi_i^{(n)}=p_i^ne^{\xi_i}\,,\quad \xi_i=p_ix_1+\frac{1}{p_i}x_{-1}+p_i^3x_3+\frac{1}{p_i^3}x_{-3}+\cdots+\xi_{i0}\,,\end{aligned}$$

and $c_{i,j}=\delta_{j,2N+1-i}\alpha_i$, $\alpha_i=\alpha_{2N+1-i}$, $p_i^2-p_ip_{2N+1-i}+p_{2N+1-i}^2=1$.

This τ-function satisfies

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_0\cdot\tau_0=\tau_1^2\,,\tag{2.13}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)\tau_1\cdot\tau_1=\tau_0\tau_2\,,\tag{2.14}$$

Proof. To impose the pseudo 3-reduction to the τ-function in Lemma 2.2, we add a constraint $p_i^3+p_{2N+1-i}^3=p_i+p_{2N+1-i}$, $p_i\neq -p_{2N+1-i}$, i.e. $p_i^2-p_ip_{2N+1-i}+p_{2N+1-i}^2=1$, and $c_{i,j}=\delta_{j,2N+1-i}\alpha_i$, $\alpha_i=\alpha_{2N+1-i}$ [**21**]. □

Lemma 2.4. *The τ-function τ_1 of the bilinear equations (2.13) and (2.14) with the pseudo 3-reduction constraint satisfies the following relation:*

$$\tau_1=\frac{1}{c}\,g_1g_2\,,\tag{2.15}$$

with

$$g_1=\mathrm{pf}\left(2\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\,,\tag{2.16}$$

$$g_2=\mathrm{pf}\left(2\alpha_i\frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\,,\tag{2.17}$$

where $\xi_i = p_i^{-1}x_{-1} + p_i x_1 + \xi_i^0$, $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 1$, $\alpha_i = \alpha_{2N+1-i}$, $c = 2^{2N}\prod_{k=1}^{2N} p_k$.

Proof. Suppose that $\alpha_i = \alpha_{2N+1-i}$ and $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 1$ are satisfied. Then we can rewrite τ_1 as follows:

$$\begin{aligned}
\tau_1 &= \det\left(\psi_{i,j}^{(1)}\right)_{1\le i,j\le 2N} = \det\left(\delta_{j,2N+1-i}\alpha_i + \frac{1}{p_i+p_j}\left(-\frac{p_i}{p_j}\right)e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i\frac{p_ip_{2N+1-i} - p_{2N+1-i}^2}{p_j(p_i-p_j)} - \frac{1}{p_i+p_j}\frac{p_i}{p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_j(p_i-p_j)} - \frac{1}{p_i+p_j}\frac{p_i}{p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_i-p_j} - \frac{2p_i}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_i-p_j} + \left(\frac{(p_i-p_j)(p_i+1)}{(p_i+p_j)(p_j-1)} - \frac{p_i-1}{p_j-1}\right)e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_i-p_j} + \frac{(p_i-p_j)(p_i+1)}{(p_i+p_j)(p_j-1)}e^{\xi_i+\xi_j} - \frac{p_i-1}{p_j-1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{\prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}}{c}\det\left(c_{i,j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j} - \frac{p_i-1}{p_i+1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N},
\end{aligned}$$

where $c_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}$ and $c = 2^{2N}\prod_{k=1}^{2N} p_k$. Using the formula (B.1), we obtain

$$\tau_1 = \frac{\prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}}{c}\begin{vmatrix} \Psi_{1,1} & \Psi_{1,2} & \cdots & \Psi_{1,2N} & \frac{p_1-1}{p_1+1}e^{\xi_1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \Psi_{2N,1} & \Psi_{2N,2} & \cdots & \Psi_{2N,2N} & \frac{p_{2N}-1}{p_{2N}+1}e^{\xi_{2N}} \\ e^{\xi_1} & e^{\xi_2} & \cdots & e^{\xi_{2N}} & 1 \end{vmatrix},$$

where $\Psi_{i,j} = c_{i,j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}$. Then we note

$$c_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j} = -2\delta_{i,2N+1-j}\,\alpha_{2N+1-i}\frac{(p_i-1)(p_j-1)}{p_j-p_i} = -c_{j,i}\,.$$

Introducing $c_i = 2\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}$, we can write as $c_{i,j} = \delta_{j,2N+1-i}c_i$ and $c_i = -c_{2N+1-i}$. Note that the $2N\times 2N$ matrix $(\Psi_{i,j})_{1\le i,j\le 2N}$ is skew-symmetric. Thus we can use the formula (D.7):

$$\begin{aligned}\tau_1 &= \frac{\prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}}{c}\,\mathrm{pf}(\Psi_{i,j})_{1\le i,j\le 2N}\,\mathrm{pf}\left(\Psi_{ij}-\frac{p_i-1}{p_i+1}e^{\xi_i+\xi_j}+\frac{p_j-1}{p_j+1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \frac{\prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}}{c}\,\mathrm{pf}\left(2\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&\quad\times\mathrm{pf}\left(2\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{(p_i-p_j)(p_i-1)(p_j-1)}{(p_i+p_j)(p_i+1)(p_j+1)}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \frac{1}{c}\,\mathrm{pf}\left(2\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&\qquad\times\mathrm{pf}\left(2\alpha_i\frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \frac{1}{c}g_1g_2\,,\end{aligned}$$

where $c=2^{2N}\prod_{k=1}^{2N}p_k$. □

LEMMA 2.5. *The τ-function τ_0 of the bilinear equations (2.13) and (2.14) with the pseudo 3-reduction constraint satisfies the relation*

$$\tau_0=\frac{1}{c}\left(g_1g_2-D_{x_1}g_1\cdot g_2\right), \tag{2.18}$$

with

$$g_1=\mathrm{pf}\left(2\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N},$$

$$g_2=\mathrm{pf}\left(2\alpha_i\frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N},$$

where $\xi_i=p_i^{-1}x_{-1}+p_ix_1+\xi_i^0$, $p_i^2-p_ip_{2N+1-i}+p_{2N+1-i}^2=1$, $\alpha_i=\alpha_{2N+1-i}$, $c=2^{2N}\prod_{k=1}^{2N}p_k$.

PROOF. Suppose that $\alpha_i=\alpha_{2N+1-i}$ and $p_i^2-p_ip_{2N+1-i}+p_{2N+1-i}^2=1$ are satisfied. Then we can rewrite τ_0 as follows:

$$\begin{aligned}\tau_0 &= \det\left(\psi_{i,j}^{(0)}\right)_{1\le i,j\le 2N} = \det\left(\delta_{j,2N+1-i}\alpha_i+\frac{1}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i\frac{p_ip_{2N+1-i}-p_{2N+1-i}^2}{p_j(p_i-p_j)}+\frac{1}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_j(p_i-p_j)}+\frac{1}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_i-p_j}+\frac{2p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_i-p_j}\right.\\
&\qquad\left.+\left(\frac{(p_i-p_j)(p_i+1)}{(p_i+p_j)(p_j-1)}-\frac{p_i+1}{p_j-1}+\frac{2p_j}{p_j-1}\right)e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N},\end{aligned}$$

where $c = 2^{2N}\prod_{k=1}^{2N} p_k$. Using the formula (B.1), we can rewrite τ_0 as follows:

$$\tau_0 = \frac{1}{c}\begin{vmatrix} \Phi_{1,1} & \Phi_{1,2} & \cdots & \Phi_{1,2N} & -2e^{\xi_1} & (p_1+1)e^{\xi_1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \Phi_{2N,1} & \Phi_{2N,2} & \cdots & \Phi_{2N,2N} & -2e^{\xi_{2N}} & (p_{2N}+1)e^{\xi_{2N}} \\ \frac{p_1}{p_1-1}e^{\xi_1} & \frac{p_2}{p_2-1}e^{\xi_2} & \cdots & \frac{p_{2N}}{p_{2N}-1}e^{\xi_{2N}} & 1 & 0 \\ \frac{1}{p_1-1}e^{\xi_1} & \frac{1}{p_2-1}e^{\xi_2} & \cdots & \frac{1}{p_{2N}-1}e^{\xi_{2N}} & 0 & 1 \end{vmatrix}$$
$$= \frac{1}{c}\begin{vmatrix} \Phi_{1,1} & \Phi_{1,2} & \cdots & \Phi_{1,2N} & (p_1-1)e^{\xi_1} & (p_1+1)e^{\xi_1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \Phi_{2N,1} & \Phi_{2N,2} & \cdots & \Phi_{2N,2N} & (p_{2N}-1)e^{\xi_{2N}} & (p_{2N}+1)e^{\xi_{2N}} \\ \frac{p_1}{p_1-1}e^{\xi_1} & \frac{p_2}{p_2-1}e^{\xi_2} & \cdots & \frac{p_{2N}}{p_{2N}-1}e^{\xi_{2N}} & 1 & 0 \\ \frac{1}{p_1-1}e^{\xi_1} & \frac{1}{p_2-1}e^{\xi_2} & \cdots & \frac{1}{p_{2N}-1}e^{\xi_{2N}} & 1 & 1 \end{vmatrix},$$

where $\Phi_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i\frac{p_i^2-1}{p_i-p_j} + \frac{(p_i-p_j)(p_i+1)}{(p_i+p_j)(p_j-1)}e^{\xi_i+\xi_j}$. Then we can further simplify as follows:

$$\tau_0 = \frac{\prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}}{c}\begin{vmatrix} \Psi_{1,1} & \Psi_{1,2} & \cdots & \Psi_{1,2N} & \frac{p_1-1}{p_1+1}e^{\xi_1} & e^{\xi_1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \Psi_{2N,1} & \Psi_{2N,2} & \cdots & \Psi_{2N,2N} & \frac{p_{2N}-1}{p_{2N}+1}e^{\xi_{2N}} & e^{\xi_{2N}} \\ p_1e^{\xi_1} & p_2e^{\xi_2} & \cdots & p_{2N}e^{\xi_{2N}} & 1 & 0 \\ e^{\xi_1} & e^{\xi_2} & \cdots & e^{\xi_{2N}} & 1 & 1 \end{vmatrix},$$

where $\Psi_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i\frac{(p_i-1)(p_j-1)}{p_i-p_j} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}$. Note that the $2N\times 2N$ matrix

$(\Psi_{i,j})_{1\le i,j\le 2N}$ is skew-symmetric. Thus we can use the formula (E.1):

$$
\begin{aligned}
\tau_0 = \frac{\prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}}{c}
&\left(
\left|\begin{array}{cccccc}
\Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & e^{\xi_1} & e^{\xi_1} \\
 & \Psi_{2,3} & \cdots & \Psi_{2,2N} & e^{\xi_2} & e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \Psi_{2N-1,2N} & e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\
 & & & & e^{\xi_{2N}} & e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right|
\right. \\
&\times
\left|\begin{array}{cccccc}
\Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & \frac{p_1-1}{p_1+1}e^{\xi_1} & p_1e^{\xi_1} \\
 & \Psi_{2,3} & \cdots & \Psi_{2,2N} & \frac{p_2-1}{p_2+1}e^{\xi_2} & p_2e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \Psi_{2N-1,2N} & \frac{p_{2N-1}-1}{p_{2N-1}+1}e^{\xi_{2N-1}} & p_{2N-1}e^{\xi_{2N-1}} \\
 & & & & \frac{p_{2N}-1}{p_{2N}+1}e^{\xi_{2N}} & p_{2N}e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right| \\
&-
\left|\begin{array}{cccccc}
\Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & e^{\xi_1} & p_1e^{\xi_1} \\
 & \Psi_{2,3} & \cdots & \Psi_{2,2N} & e^{\xi_2} & p_2e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \Psi_{2N-1,2N} & e^{\xi_{2N-1}} & p_{2N-1}e^{\xi_{2N-1}} \\
 & & & & e^{\xi_{2N}} & p_{2N}e^{\xi_{2N}} \\
 & & & & & 0
\end{array}\right| \\
&\left.\times
\left|\begin{array}{cccccc}
\Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & \frac{p_1-1}{p_1+1}e^{\xi_1} & e^{\xi_1} \\
 & \Psi_{2,3} & \cdots & \Psi_{2,2N} & \frac{p_2-1}{p_2+1}e^{\xi_2} & e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \Psi_{2N-1,2N} & \frac{p_{2N-1}-1}{p_{2N-1}+1}e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\
 & & & & \frac{p_{2N}-1}{p_{2N}+1}e^{\xi_{2N}} & e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right|
\right)
\end{aligned}
$$

Let

$$
g_1 = \mathrm{pf}\,(\Psi_{i,j})_{1\le i,j\le 2N} =
\left|\begin{array}{cccccc}
\Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & e^{\xi_1} & e^{\xi_1} \\
 & \Psi_{2,3} & \cdots & \Psi_{2,2N} & e^{\xi_2} & e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \Psi_{2N-1,2N} & e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\
 & & & & e^{\xi_{2N}} & e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right|,
$$

$$
\begin{aligned}
g_2 &= \mathrm{pf}\left(2\alpha_i\frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}\,\mathrm{pf}\left(\Psi_{ij} - \frac{p_i-1}{p_i+1}e^{\xi_i+\xi_j} + \frac{p_j-1}{p_j+1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}\times
\left|\begin{array}{cccccc}
\Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & \frac{p_1-1}{p_1+1}e^{\xi_1} & e^{\xi_1} \\
 & \Psi_{2,3} & \cdots & \Psi_{2,2N} & \frac{p_2-1}{p_2+1}e^{\xi_2} & e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \Psi_{2N-1,2N} & \frac{p_{2N-1}-1}{p_{2N-1}+1}e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\
 & & & & \frac{p_{2N}-1}{p_{2N}+1}e^{\xi_{2N}} & e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right|.
\end{aligned}
$$

Then

$$\partial_{x_1} g_1 = \left| \begin{array}{ccccc} \Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & e^{\xi_1} & p_1 e^{\xi_1} \\ & \Psi_{2,3} & \cdots & \Psi_{2,2N} & e^{\xi_2} & p_2 e^{\xi_2} \\ & & \ddots & \vdots & \vdots & \vdots \\ & & & \Psi_{2N-1,2N} & e^{\xi_{2N-1}} & p_{2N-1} e^{\xi_{2N-1}} \\ & & & & e^{\xi_{2N}} & p_{2N} e^{\xi_{2N}} \\ & & & & & 0 \end{array} \right|,$$

$$\begin{aligned} &(\partial_{x_1}+1)g_2 = (\partial_{x_1}+1)\mathrm{pf}\left(2\alpha_i \frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\ &= \mathrm{pf}\left(2\alpha_i \frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j} + (p_i-p_j)e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\ &= \prod_{k=1}^{2N}\frac{p_k+1}{p_k-1}\,\mathrm{pf}\left(\Psi_{ij} - \frac{p_i-1}{p_i+1}p_j e^{\xi_i+\xi_j} + p_i\frac{p_j-1}{p_j+1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\ &= \prod_{k=1}^{2N}\frac{p_k+1}{p_k-1} \times \left| \begin{array}{ccccc} \Psi_{1,2} & \Psi_{1,3} & \cdots & \Psi_{1,2N} & \frac{p_1-1}{p_1+1}e^{\xi_1} & p_1 e^{\xi_1} \\ & \Psi_{2,3} & \cdots & \Psi_{2,2N} & \frac{p_2-1}{p_2+1}e^{\xi_2} & p_2 e^{\xi_2} \\ & & \ddots & \vdots & \vdots & \vdots \\ & & & \Psi_{2N-1,2N} & \frac{p_{2N-1}-1}{p_{2N-1}+1}e^{\xi_{2N-1}} & p_{2N-1} e^{\xi_{2N-1}} \\ & & & & \frac{p_{2N}-1}{p_{2N}+1}e^{\xi_{2N}} & p_{2N} e^{\xi_{2N}} \\ & & & & & 1 \end{array} \right|. \end{aligned}$$

Thus τ_0, g_1 and g_2 satisfy the relation (2.18). □

LEMMA 2.6. *The τ-function τ_2 of the bilinear equations (2.13) and (2.14) with the pseudo 3-reduction constraint satisfies the relation*

$$\tau_2 = \frac{1}{c}\left(g_1 g_2 - D_{x_{-1}} g_1 \cdot g_2\right), \tag{2.19}$$

with

$$\begin{aligned} g_1 &= \mathrm{pf}\left(2\alpha_i \frac{(p_i-1)(p_j-1)}{p_i-p_j}\delta_{j,2N+1-i} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}, \\ g_2 &= \mathrm{pf}\left(2\alpha_i \frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i} + \frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}, \end{aligned}$$

where $\xi_i = p_i^{-1}x_{-1} + p_i x_1 + \xi_i^0$, $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 1$, $\alpha_i = \alpha_{2N+1-i}$, $c = 2^{2N}\prod_{k=1}^{2N} p_k$.

PROOF. Suppose that $\alpha_i = \alpha_{2N+1-i}$ and $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 1$ are satisfied. Then we can rewrite τ_2 as follows:

$$\begin{aligned}
\tau_2 &= \det\left(\psi_{i,j}^{(2)}\right)_{1\le i,j\le 2N} \\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i \frac{p_{2N+1-i}(p_i - p_{2N+1-i})}{p_j(p_i-p_j)} + \frac{1}{p_i+p_j}\frac{p_i^2}{p_j^2} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i \frac{p_i p_{2N+1-i} - p_{2N+1-i}^2}{p_j(p_i-p_j)} + \frac{1}{p_i+p_j}\frac{p_i^2}{p_j^2} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \det\left(\delta_{j,2N+1-i}\,\alpha_i \frac{p_i^2-1}{p_j(p_i-p_j)} + \frac{1}{p_i+p_j}\frac{p_i^2}{p_j^2} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i \frac{p_i^2-1}{p_i-p_j} + \frac{2p_i^2}{p_i+p_j}\frac{1}{p_j} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i \frac{p_j}{p_i}\frac{p_i^2-1}{p_i-p_j} + \frac{2p_i}{p_i+p_j} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\left(2\delta_{j,2N+1-i}\,\alpha_i \frac{p_i^{-2}-1}{p_i^{-1}-p_j^{-1}} + \frac{2p_i}{p_i+p_j} e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \frac{1}{c}\det\Bigg(2\delta_{j,2N+1-i}\,\alpha_i \frac{p_i^{-2}-1}{p_i^{-1}-p_j^{-1}} \\
&\qquad + \left(\frac{(p_i^{-1}-p_j^{-1})(p_i^{-1}+1)}{(p_i^{-1}+p_j^{-1})(p_j^{-1}-1)} - \frac{p_i^{-1}+1}{p_j^{-1}-1} + \frac{2p_j^{-1}}{p_j^{-1}-1}\right) e^{\xi_i+\xi_j}\Bigg)_{1\le i,j\le 2N},
\end{aligned}$$

where $c = 2^{2N}\prod_{k=1}^{2N} p_k$. Using the formula (B.1), we can rewrite τ_2 as follows:

$$\begin{aligned}
\tau_2 &= \frac{1}{c}\begin{vmatrix}
\tilde\Phi_{1,1} & \tilde\Phi_{1,2} & \cdots & \tilde\Phi_{1,2N} & -2e^{\xi_1} & (p_1^{-1}+1)e^{\xi_1} \\
\vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\
\tilde\Phi_{2N,1} & \tilde\Phi_{2N,2} & \cdots & \tilde\Phi_{2N,2N} & -2e^{\xi_{2N}} & (p_{2N}^{-1}+1)e^{\xi_{2N}} \\
\frac{p_1^{-1}}{p_1^{-1}-1}e^{\xi_1} & \frac{p_2^{-1}}{p_2^{-1}-1}e^{\xi_2} & \cdots & \frac{p_{2N}^{-1}}{p_{2N}^{-1}-1}e^{\xi_{2N}} & 1 & 0 \\
\frac{1}{p_1^{-1}-1}e^{\xi_1} & \frac{1}{p_2^{-1}-1}e^{\xi_2} & \cdots & \frac{1}{p_{2N}^{-1}-1}e^{\xi_{2N}} & 0 & 1
\end{vmatrix} \\
&= \frac{1}{c}\begin{vmatrix}
\tilde\Phi_{1,1} & \tilde\Phi_{1,2} & \cdots & \tilde\Phi_{1,2N} & (p_1^{-1}-1)e^{\xi_1} & (p_1^{-1}+1)e^{\xi_1} \\
\vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\
\tilde\Phi_{2N,1} & \tilde\Phi_{2N,2} & \cdots & \tilde\Phi_{2N,2N} & (p_{2N}^{-1}-1)e^{\xi_{2N}} & (p_{2N}^{-1}+1)e^{\xi_{2N}} \\
\frac{p_1^{-1}}{p_1^{-1}-1}e^{\xi_1} & \frac{p_2^{-1}}{p_2^{-1}-1}e^{\xi_2} & \cdots & \frac{p_{2N}^{-1}}{p_{2N}^{-1}-1}e^{\xi_{2N}} & 1 & 0 \\
\frac{1}{p_1^{-1}-1}e^{\xi_1} & \frac{1}{p_2^{-1}-1}e^{\xi_2} & \cdots & \frac{1}{p_{2N}^{-1}-1}e^{\xi_{2N}} & 1 & 1
\end{vmatrix},
\end{aligned}$$

where $\tilde{\Phi}_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i \frac{p_i^{-2}-1}{p_i^{-1}-p_j^{-1}} + \frac{(p_i^{-1}-p_j^{-1})(p_i^{-1}+1)}{(p_i^{-1}+p_j^{-1})(p_j^{-1}-1)} e^{\xi_i+\xi_j}$. Then we can further simplify as follows:

$$\tau_2 = \frac{\prod_{k=1}^{2N} \frac{p_k^{-1}+1}{p_k^{-1}-1}}{c} \begin{vmatrix} \tilde{\Psi}_{1,1} & \tilde{\Psi}_{1,2} & \cdots & \tilde{\Psi}_{1,2N} & \frac{p_1^{-1}-1}{p_1^{-1}+1} e^{\xi_1} & e^{\xi_1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \tilde{\Psi}_{2N,1} & \tilde{\Psi}_{2N,2} & \cdots & \tilde{\Psi}_{2N,2N} & \frac{p_{2N}^{-1}-1}{p_{2N}^{-1}+1} e^{\xi_{2N}} & e^{\xi_{2N}} \\ p_1^{-1} e^{\xi_1} & p_2^{-1} e^{\xi_2} & \cdots & p_{2N}^{-1} e^{\xi_{2N}} & 1 & 0 \\ e^{\xi_1} & e^{\xi_2} & \cdots & e^{\xi_{2N}} & 1 & 1 \end{vmatrix},$$

where $\tilde{\Psi}_{i,j} = 2\delta_{j,2N+1-i}\,\alpha_i \frac{(p_i^{-1}-1)(p_j^{-1}-1)}{p_i^{-1}-p_j^{-1}} + \frac{p_i^{-1}-p_j^{-1}}{p_i^{-1}+p_j^{-1}} e^{\xi_i+\xi_j}$. Note that the $2N \times 2N$ matrix $(\tilde{\Psi}_{i,j})_{1 \le i,j \le 2N}$ is skew-symmetric and $\tilde{\Psi}_{i,j} = -\Psi_{i,j}$. Thus we can use the formula (E.1):

$$\tau_2 = \frac{\prod_{k=1}^{2N} \frac{p_k^{-1}+1}{p_k^{-1}-1}}{c} \left(\left. \begin{array}{ccccccc} | & \tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & e^{\xi_1} & e^{\xi_1} \\ & & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & e^{\xi_2} & e^{\xi_2} \\ & & & \ddots & \vdots & \vdots & \vdots \\ & & & & \tilde{\Psi}_{2N-1,2N} & e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\ & & & & & e^{\xi_{2N}} & e^{\xi_{2N}} \\ & & & & & & 1 \end{array} \right| \right.$$

$$\left. \times \left. \begin{array}{ccccccc} | & \tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & \frac{p_1^{-1}-1}{p_1^{-1}+1} e^{\xi_1} & p_1^{-1} e^{\xi_1} \\ & & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & \frac{p_2^{-1}-1}{p_2^{-1}+1} e^{\xi_2} & p_2^{-1} e^{\xi_2} \\ & & & \ddots & \vdots & \vdots & \vdots \\ & & & & \tilde{\Psi}_{2N-1,2N} & \frac{p_{2N-1}^{-1}-1}{p_{2N-1}^{-1}+1} e^{\xi_{2N-1}} & p_{2N-1}^{-1} e^{\xi_{2N-1}} \\ & & & & & \frac{p_{2N}^{-1}-1}{p_{2N}^{-1}+1} e^{\xi_{2N}} & p_{2N}^{-1} e^{\xi_{2N}} \\ & & & & & & 1 \end{array} \right| \right.$$

$$\left. - \left. \begin{array}{ccccccc} | & \tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & e^{\xi_1} & p_1^{-1} e^{\xi_1} \\ & & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & e^{\xi_2} & p_2^{-1} e^{\xi_2} \\ & & & \ddots & \vdots & \vdots & \vdots \\ & & & & \tilde{\Psi}_{2N-1,2N} & e^{\xi_{2N-1}} & p_{2N-1}^{-1} e^{\xi_{2N-1}} \\ & & & & & e^{\xi_{2N}} & p_{2N}^{-1} e^{\xi_{2N}} \\ & & & & & & 0 \end{array} \right| \right.$$

$$\left. \times \left. \begin{array}{ccccccc} | & \tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & \frac{p_1^{-1}-1}{p_1^{-1}+1} e^{\xi_1} & e^{\xi_1} \\ & & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & \frac{p_2^{-1}-1}{p_2^{-1}+1} e^{\xi_2} & e^{\xi_2} \\ & & & \ddots & \vdots & \vdots & \vdots \\ & & & & \tilde{\Psi}_{2N-1,2N} & \frac{p_{2N-1}^{-1}-1}{p_{2N-1}^{-1}+1} e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\ & & & & & \frac{p_{2N}^{-1}-1}{p_{2N}^{-1}+1} e^{\xi_{2N}} & e^{\xi_{2N}} \\ & & & & & & 1 \end{array} \right| \right).$$

Let

$$
\begin{aligned}
g_1 &= \mathrm{pf}\left(\Psi_{i,j}\right)_{1\le i,j\le 2N} = \mathrm{pf}\left(\tilde{\Psi}_{i,j}\right)_{1\le i,j\le 2N} \\
&= \left|\begin{array}{cccccc}
\tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & e^{\xi_1} & e^{\xi_1} \\
 & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & e^{\xi_2} & e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \tilde{\Psi}_{2N-1,2N} & e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\
 & & & & e^{\xi_{2N}} & e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right|,
\end{aligned}
$$

$$
\begin{aligned}
g_2 &= \mathrm{pf}\left(2\alpha_i\frac{(p_i+1)(p_j+1)}{p_i-p_j}\delta_{j,2N+1-i}+\frac{p_i-p_j}{p_i+p_j}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \mathrm{pf}\left(2\alpha_i\frac{(p_i^{-1}+1)(p_j^{-1}+1)}{p_i^{-1}-p_j^{-1}}\delta_{j,2N+1-i}+\frac{p_i^{-1}-p_j^{-1}}{p_i^{-1}+p_j^{-1}}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \prod_{k=1}^{2N}\frac{p_k^{-1}+1}{p_k^{-1}-1}\,\mathrm{pf}\left(\tilde{\Psi}_{ij}-\frac{p_i^{-1}-1}{p_i^{-1}+1}e^{\xi_i+\xi_j}+\frac{p_j^{-1}-1}{p_j^{-1}+1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N} \\
&= \prod_{k=1}^{2N}\frac{p_k^{-1}+1}{p_k^{-1}-1}\times\left|\begin{array}{cccccc}
\tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & \frac{p_1^{-1}-1}{p_1^{-1}+1}e^{\xi_1} & e^{\xi_1} \\
 & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & \frac{p_2^{-1}-1}{p_2^{-1}+1}e^{\xi_2} & e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \tilde{\Psi}_{2N-1,2N} & \frac{p_{2N-1}^{-1}-1}{p_{2N-1}^{-1}+1}e^{\xi_{2N-1}} & e^{\xi_{2N-1}} \\
 & & & & \frac{p_{2N}^{-1}-1}{p_{2N}^{-1}+1}e^{\xi_{2N}} & e^{\xi_{2N}} \\
 & & & & & 1
\end{array}\right|.
\end{aligned}
$$

Then

$$
\partial_{x_{-1}}g_1 = \left|\begin{array}{cccccc}
\tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & e^{\xi_1} & p_1^{-1}e^{\xi_1} \\
 & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & e^{\xi_2} & p_2^{-1}e^{\xi_2} \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \tilde{\Psi}_{2N-1,2N} & e^{\xi_{2N-1}} & p_{2N-1}^{-1}e^{\xi_{2N-1}} \\
 & & & & e^{\xi_{2N}} & p_{2N}^{-1}e^{\xi_{2N}} \\
 & & & & & 0
\end{array}\right|,
$$

$$
\begin{aligned}
&(\partial_{x_{-1}}+1)g_2\\
&=(\partial_{x_{-1}}+1)\mathrm{pf}\left(2\alpha_i\frac{(p_i^{-1}+1)(p_j^{-1}+1)}{p_i^{-1}-p_j^{-1}}\delta_{j,2N+1-i}+\frac{p_i^{-1}-p_j^{-1}}{p_i^{-1}+p_j^{-1}}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&=\mathrm{pf}\left(2\alpha_i\frac{(p_i^{-1}+1)(p_j^{-1}+1)}{p_i^{-1}-p_j^{-1}}\delta_{j,2N+1-i}+\frac{p_i^{-1}-p_j^{-1}}{p_i^{-1}+p_j^{-1}}e^{\xi_i+\xi_j}+(p_i^{-1}-p_j^{-1})e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&=\prod_{k=1}^{2N}\frac{p_k^{-1}+1}{p_k^{-1}-1}\,\mathrm{pf}\left(\tilde{\Psi}_{ij}-\frac{p_i^{-1}-1}{p_i^{-1}+1}p_j^{-1}e^{\xi_i+\xi_j}+p_i^{-1}\frac{p_j^{-1}-1}{p_j^{-1}+1}e^{\xi_i+\xi_j}\right)_{1\le i,j\le 2N}\\
&=\prod_{k=1}^{2N}\frac{p_k^{-1}+1}{p_k^{-1}-1}\times
\begin{array}{|cccccc|}
\tilde{\Psi}_{1,2} & \tilde{\Psi}_{1,3} & \cdots & \tilde{\Psi}_{1,2N} & \frac{p_1^{-1}-1}{p_1^{-1}+1}e^{\xi_1} & p_1^{-1}e^{\xi_1}\\
 & \tilde{\Psi}_{2,3} & \cdots & \tilde{\Psi}_{2,2N} & \frac{p_2^{-1}-1}{p_2^{-1}+1}e^{\xi_2} & p_2^{-1}e^{\xi_2}\\
 & & \ddots & \vdots & \vdots & \vdots\\
 & & & \tilde{\Psi}_{2N-1,2N} & \frac{p_{2N-1}^{-1}-1}{p_{2N-1}^{-1}+1}e^{\xi_{2N-1}} & p_{2N-1}^{-1}e^{\xi_{2N-1}}\\
 & & & & \frac{p_{2N}^{-1}-1}{p_{2N}^{-1}+1}e^{\xi_{2N}} & p_{2N}^{-1}e^{\xi_{2N}}\\
 & & & & & 1
\end{array}\,.
\end{aligned}
$$

Thus τ_2, g_1 and g_2 satisfy the relation (2.19). □

Letting $F=\tau_0$, $G=\tau_1$ and $H=\tau_2$, we obtain the following equations

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)F\cdot F=G^2\,, \tag{2.20}$$

$$-\left(\frac{1}{2}D_{x_1}D_{x_{-1}}-1\right)G\cdot G=FH\,, \tag{2.21}$$

$$cG=g_1g_2\,, \tag{2.22}$$

$$cF=g_1g_2-D_{x_1}g_1\cdot g_2\,, \tag{2.23}$$

$$cH=g_1g_2-D_{x_{-1}}g_1\cdot g_2\,, \tag{2.24}$$

from the bilinear equations (2.13) and (2.14), and the relations between determinants and pfaffians (2.15), (2.18), (2.19).

Theorem 2.7. *The τ-functions g_1 (2.16) and g_2 (2.17) of equations (2.20), (2.21), (2.22), (2.23) and (2.24) give the N-soliton solution of the DP equation*

$$u_t+3u_x-u_{txx}+4uu_x=3u_xu_{xx}+uu_{xxx}\,, \tag{2.25}$$

through the dependent variable transformation

$$u=-\left(\ln\frac{g_1}{g_2}\right)_{x_{-1}}\,,$$

and the hodograph (reciprocal) transformation

$$\begin{cases} x=x_1+\int_{-\infty}^{x_{-1}}u(x_1,x'_{-1})dx'_{-1}\\ \quad =x_1-\ln\frac{g_1}{g_2}\,,\\ t=x_{-1}\,. \end{cases} \tag{2.26}$$

Proof. From (2.22), (2.23) and (2.24), we have the relations

$$-\left(\ln\frac{g_1}{g_2}\right)_{x_1}=\frac{F}{G}-1\,, \tag{2.27}$$

$$-\left(\ln\frac{g_1}{g_2}\right)_{x_{-1}}=\frac{H}{G}-1\,. \tag{2.28}$$

Let

$$\rho=\frac{G}{F}\,,\quad u=-\left(\ln\frac{g_1}{g_2}\right)_{x_{-1}}\,. \tag{2.29}$$

Differentiating (2.27) with respect to x_{-1}, we obtain

$$u_{x_1}=\left(\frac{1}{\rho}\right)_{x_{-1}}\,. \tag{2.30}$$

This is rewritten as

$$(\ln\rho)_{x_{-1}}=-\rho u_{x_1}\,. \tag{2.31}$$

Equation (2.28) leads to

$$\frac{H}{G}=1+u\,. \tag{2.32}$$

The bilinear equations (2.20) and (2.21) are written as

$$-\,(\ln F)_{x_1x_{-1}}+1=\rho^2\,, \tag{2.33}$$

$$-\,(\ln G)_{x_1x_{-1}}+1=\frac{1}{\rho}(1+u)\,. \tag{2.34}$$

Subtracting (2.33) from (2.34), we obtain

$$-\,(\ln\rho)_{x_1x_{-1}}=\frac{1}{\rho}(1+u)-\rho^2\,, \tag{2.35}$$

which leads to

$$\rho^3=1+u+\rho(\ln\rho)_{x_1x_{-1}}\,. \tag{2.36}$$

Using (2.31), it becomes

$$\rho^3=1+u-\rho(\rho u_{x_1})_{x_1}\,. \tag{2.37}$$

Let us consider the hodograph (reciprocal) transformation

$$\left\{\begin{array}{l} x=x_1+\int_{-\infty}^{x_{-1}}u(x_1,x'_{-1})dx'_{-1}\\ \quad =x_1-\ln\frac{g_1}{g_2}\,,\\ t=x_{-1}\,. \end{array}\right. \tag{2.38}$$

This yields

$$\left\{\begin{array}{l} \frac{\partial x}{\partial x_1}=1-\left(\ln\frac{g_1}{g_2}\right)_{x_1}=\rho^{-1}\,,\\ \frac{\partial x}{\partial x_{-1}}=-\left(\ln\frac{g_1}{g_2}\right)_{x_{-1}}=u\,, \end{array}\right. \tag{2.39}$$

and

$$\left\{\begin{array}{l} \partial_{x_1}=\frac{1}{\rho}\partial_x\,,\\ \partial_{x_{-1}}=\partial_t+u\partial_x\,. \end{array}\right. \tag{2.40}$$

Applying the hodograph (reciprocal) transformation to (2.31) and (2.37), we obtain

$$\left\{\begin{array}{l} (\partial_t + u\partial_x)\ln\rho = -u_x\,, \\ \rho^3 = 1 + u - u_{xx}\,. \end{array}\right. \tag{2.41}$$

This is equivalent to

$$(\partial_t + u\partial_x)\ln(1 + u - u_{xx}) = -3u_x\,, \tag{2.42}$$

which can be written as

$$(\partial_t + u\partial_x)(1 + u - u_{xx}) = -3u_x(1 + u - u_{xx})\,. \tag{2.43}$$

This is nothing but the DP equation (2.25). □

Remark 2.8. Applying the scale transformation $u \to \frac{1}{\kappa^3}u$, $t \to \kappa^3 t$ to (2.25), we obtain the DP equation (1.1).

Let $k_i = p_i + p_{2N+1-i}$. From $p_i^2 - p_i p_{2N+1-i} + p_{2N+1-i}^2 = 1$, we obtain $p_i = \frac{1}{6}(3k_i + \sqrt{3(4-k_i^2)})$, $p_{2N+1-i} = \frac{1}{6}(3k_i - \sqrt{3(4-k_i^2)})$, $p_i p_{2N+1-i} = \frac{k_i^2-1}{3}$ and $\frac{1}{pi} + \frac{1}{p_{2N+1-i}} = \frac{3k_i}{k_i^2-1}$. Thus

$$\xi_i + \xi_{2N+1-i} = k_i x_1 + \frac{3k_i}{k_i^2-1}x_{-1} + \xi_{i0} + \xi_{2N+1-i0}\,.$$

In the pfaffian solution, all phase functions can be expressed by the summation of $\xi_i + \xi_{2N+1-i}$. So the phase functions can be expressed by the parameters $\{k_i\}$ $(i = 1, 2, \cdots, N)$. Each coefficient of exponential functions can be normalized to 1 after absorption into phase constants or can be rewritten by the parameters $\{k_i\}$. Thus it is possible to rewrite the above τ-function by using the parameters $\{k_i\}$ instead of $\{p_i\}$.

Example 2.9. Soliton Solutions
For $N = 1$,

$$\begin{aligned} g_1 &= 2\alpha_1\frac{(p_1-1)(p_2-1)}{p_1-p_2} + \frac{p_1-p_2}{p_1+p_2}e^{\xi_1+\xi_2} \\ &= \frac{(p_1-1)(p_2-1)}{p_1-p_2}\left(2\alpha_1 + \frac{(p_1-p_2)^2}{(p_1+p_2)(p_1-1)(p_2-1)}e^{\xi_1+\xi_2}\right)\,. \end{aligned}$$

$$\begin{aligned} g_2 &= 2\alpha_1\frac{(p_1+1)(p_2+1)}{p_1-p_2} + \frac{p_1-p_2}{p_1+p_2}e^{\xi_1+\xi_2} \\ &= \frac{(p_1+1)(p_2+1)}{p_1-p_2}\left(2\alpha_1 + \frac{(p_1-p_2)^2}{(p_1+p_2)(p_1+1)(p_2+1)}e^{\xi_1+\xi_2}\right)\,. \end{aligned}$$

Letting $\alpha_1 = \frac{1}{2}$ and $e^{\gamma_1} = \frac{(p_1-p_2)^2}{(p_1+p_2)}\frac{1}{\sqrt{(p_1-1)(p_2-1)(p_1+1)(p_2+1)}}$, the τ-functions can be rewritten as

$$g_1 = 1 + e^{\xi_1+\xi_2+\phi_1+\gamma_1}\,, \quad g_2 = 1 + e^{\xi_1+\xi_2-\phi_1+\gamma_1} \tag{2.44}$$

where

$$\xi_1 + \xi_2 = k_1 x_1 + \frac{3k_1}{k_1^2-1}x_{-1} + \xi_{10} + \xi_{20}\,,$$

and

$$e^{\phi_1}=\sqrt{\frac{(p_1+1)(p_2+1)}{(p_1-1)(p_2-1)}}=\sqrt{\frac{k_1^2+3k_1+2}{k_1^2-3k_1+2}}=\sqrt{\frac{(k_1+2)(k_1+1)}{(k_1-2)(k_1-1)}}\,.$$

Here γ_1 can be absorbed into a phase constant.
For $N=2$,

$$\begin{aligned}
g_1&=\frac{p_1-p_2}{p_1+p_2}e^{\xi_1+\xi_2}\frac{p_3-p_4}{p_3+p_4}e^{\xi_3+\xi_4}-\frac{p_1-p_3}{p_1+p_3}e^{\xi_1+\xi_3}\frac{p_2-p_4}{p_2+p_4}e^{\xi_2+\xi_4}\\
&+\left(2\alpha_1\frac{(p_1-1)(p_4-1)}{p_1-p_4}+\frac{p_1-p_4}{p_1+p_4}e^{\xi_1+\xi_4}\right)\left(2\alpha_2\frac{(p_2-1)(p_3-1)}{p_2-p_3}+\frac{p_2-p_3}{p_2+p_3}e^{\xi_2+\xi_3}\right)\\
&=\frac{(p_1-1)(p_4-1)}{p_1-p_4}\cdot\frac{(p_2-1)(p_3-1)}{p_2-p_3}\left(2\alpha_1\cdot2\alpha_2+2\alpha_2\frac{(p_1-p_4)^2}{(p_1+p_4)(p_1-1)(p_4-1)}e^{\xi_1+\xi_4}\right.\\
&+2\alpha_1\frac{(p_2-p_3)^2}{(p_2+p_3)(p_2-1)(p_3-1)}e^{\xi_2+\xi_3}+\frac{p_1-p_4}{(p_1-1)(p_4-1)}\cdot\frac{p_2-p_3}{(p_2-1)(p_3-1)}\\
&\left.\times\left(\frac{p_1-p_2}{p_1+p_2}\frac{p_3-p_4}{p_3+p_4}-\frac{p_1-p_3}{p_1+p_3}\frac{p_2-p_4}{p_2+p_4}+\frac{p_1-p_4}{p_1+p_4}\frac{p_2-p_3}{p_2+p_3}\right)e^{\xi_1+\xi_2+\xi_3+\xi_4}\right)\,,
\end{aligned}$$

$$\begin{aligned}
g_2&=\frac{p_1-p_2}{p_1+p_2}e^{\xi_1+\xi_2}\frac{p_3-p_4}{p_3+p_4}e^{\xi_3+\xi_4}-\frac{p_1-p_3}{p_1+p_3}e^{\xi_1+\xi_3}\frac{p_2-p_4}{p_2+p_4}e^{\xi_2+\xi_4}\\
&+\left(2\alpha_1\frac{(p_1+1)(p_4+1)}{p_1-p_4}+\frac{p_1-p_4}{p_1+p_4}e^{\xi_1+\xi_4}\right)\left(2\alpha_2\frac{(p_2+1)(p_3+1)}{p_2-p_3}+\frac{p_2-p_3}{p_2+p_3}e^{\xi_2+\xi_3}\right)\\
&=\frac{(p_1+1)(p_4+1)}{p_1-p_4}\cdot\frac{(p_2+1)(p_3+1)}{p_2-p_3}\left(2\alpha_1\cdot2\alpha_2+2\alpha_2\frac{(p_1-p_4)^2}{(p_1+p_4)(p_1+1)(p_4+1)}e^{\xi_1+\xi_4}\right.\\
&+2\alpha_1\frac{(p_2-p_3)^2}{(p_2+p_3)(p_2+1)(p_3+1)}e^{\xi_2+\xi_3}+\frac{p_1-p_4}{(p_1+1)(p_4+1)}\cdot\frac{p_2-p_3}{(p_2+1)(p_3+1)}\\
&\left.\times\left(\frac{p_1-p_2}{p_1+p_2}\frac{p_3-p_4}{p_3+p_4}-\frac{p_1-p_3}{p_1+p_3}\frac{p_2-p_4}{p_2+p_4}+\frac{p_1-p_4}{p_1+p_4}\frac{p_2-p_3}{p_2+p_3}\right)e^{\xi_1+\xi_2+\xi_3+\xi_4}\right)\,.
\end{aligned}$$

Letting $\alpha_1=\alpha_2=\frac{1}{2}$, $e^{\gamma_1}=\frac{(p_1-p_4)^2}{(p_1+p_4)}\frac{1}{\sqrt{(p_1-1)(p_4-1)(p_1+1)(p_4+1)}}$, $e^{\gamma_2}=\frac{(p_2-p_3)^2}{(p_2+p_3)}\frac{1}{\sqrt{(p_2-1)(p_3-1)(p_2+1)(p_3+1)}}$, the above τ-functions become

$$\begin{aligned}
g_1&=1+e^{\xi_1+\xi_4+\phi_1}+e^{\xi_2+\xi_3+\phi_2}+b_{12}e^{\xi_1+\xi_2+\xi_3+\xi_4+\phi_1+\phi_2}\,,\\
g_2&=1+e^{\xi_1+\xi_4-\phi_1}+e^{\xi_2+\xi_3-\phi_2}+b_{12}e^{\xi_1+\xi_2+\xi_3+\xi_4-\phi_1-\phi_2}\,,
\end{aligned}$$

where

$$\begin{aligned}
b_{12} &= \frac{p_1-p_2}{p_1+p_2}\frac{p_3-p_4}{p_3+p_4}\frac{p_1+p_4}{p_1-p_4}\frac{p_2+p_3}{p_2-p_3} - \frac{p_1-p_3}{p_1+p_3}\frac{p_2-p_4}{p_2+p_4}\frac{p_1+p_4}{p_1-p_4}\frac{p_2+p_3}{p_2-p_3} + 1 \\
&= \frac{(k_1-k_2)^2(k_1^2-k_1k_2+k_2^2-3)}{(k_1+k_2)^2(k_1^2+k_1k_2+k_2^2-3)}\,, \\
\xi_1+\xi_4 &= k_1x_1 + \frac{3k_1}{k_1^2-1}x_{-1} + \xi_{10} + \xi_{40}\,, \quad \xi_2+\xi_3 = k_2x_1 + \frac{3k_2}{k_2^2-1}x_{-1} + \xi_{20} + \xi_{30}\,, \\
e^{\phi_1} &= \sqrt{\frac{(p_1+1)(p_4+1)}{(p_1-1)(p_4-1)}} = \sqrt{\frac{k_1^2+3k_1+2}{k_1^2-3k_1+2}} = \sqrt{\frac{(k_1+2)(k_1+1)}{(k_1-2)(k_1-1)}}\,, \\
e^{\phi_2} &= \sqrt{\frac{(p_2+1)(p_3+1)}{(p_2-1)(p_3-1)}} = \sqrt{\frac{k_2^2+3k_2+2}{k_2^2-3k_2+2}} = \sqrt{\frac{(k_2+2)(k_2+1)}{(k_2-2)(k_2-1)}}\,.
\end{aligned}$$

Here γ_1 and γ_2 were absorbed into phase constants.

The N-soliton solution of (2.25) is written in the following form:

$$g_1 = \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i(\eta_i+\phi_i) + \sum_{i<j}^{(N)} \mu_i\mu_j \ln b_{ij}\right]\,, \tag{2.45}$$

$$g_2 = \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i(\eta_i-\phi_i) + \sum_{i<j}^{(N)} \mu_i\mu_j \ln b_{ij}\right]\,,$$

$$b_{ij} = \frac{(k_i-k_j)^2(k_i^2-k_ik_j+k_j^2-3)}{(k_i+k_j)^2(k_i^2+k_ik_j+k_j^2-3)} \quad \text{for} \quad i<j\,,$$

$$\eta_i = \xi_i + \xi_{2N+1-i} = k_ix_1 + \frac{3k_i}{k_i^2-1}x_{-1} + \eta_{i0}\,, \tag{2.46}$$

$$e^{\phi_i} = \sqrt{\frac{k_i^2+3k_i+2}{k_i^2-3k_i+2}} = \sqrt{\frac{(k_i+2)(k_i+1)}{(k_i-2)(k_i-1)}}\,,$$

where $\sum\limits_{\mu=0,1}$ means the summation over all possible combinations of $\mu_i = 0$ or 1 for $i = 1, 2, \cdots, N$, and $\sum\limits_{i<j}^{(N)}$ means the summation over all possible combinations of N elements under the condition $i<j$.

Applying $u \to \frac{1}{\kappa^3}u$, $t \to \kappa^3 t$, $x_{-1} \to \kappa^3 x_{-1}$, $x_1 \to \frac{x_1}{\kappa}$, $\kappa k_i = p_i + p_{2N+1-i}$, we obtain the N-soliton solution of the DP equation (1.1).

Theorem 2.10. *The N-soliton solution of the DP equation (1.1) is given as follows:*

$$
\begin{aligned}
&u = -\left(\ln \frac{g_1}{g_2}\right)_{x_{-1}} , \\
&g_1 = \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i(\eta_i + \phi_i) + \sum_{i<j}^{(N)} \mu_i\mu_j \ln b_{ij}\right] , \\
&g_2 = \sum_{\mu=0,1} \exp\left[\sum_{i=1}^{N} \mu_i(\eta_i - \phi_i) + \sum_{i<j}^{(N)} \mu_i\mu_j \ln b_{ij}\right] , \\
&b_{ij} = \frac{(k_i - k_j)^2((k_i^2 - k_ik_j + k_j^2)\kappa^2 - 3)}{(k_i + k_j)^2((k_i^2 + k_ik_j + k_j^2)\kappa^2 - 3)} \quad \text{for} \quad i < j , \\
&\eta_i = \xi_i + \xi_{2N+1-i} = k_ix_1 + \frac{3k_i\kappa^4}{\kappa^2k_i^2 - 1}x_{-1} + \eta_{i0} , \\
&e^{\phi_i} = \sqrt{\frac{\kappa^2k_i^2 + 3\kappa k_i + 2}{\kappa^2k_i^2 - 3\kappa k_i + 2}} = \sqrt{\frac{(\kappa k_i + 2)(\kappa k_i + 1)}{(\kappa k_i - 2)(\kappa k_i - 1)}} ,
\end{aligned}
$$

and the hodograph (reciprocal) transformation

$$
\left\{
\begin{array}{l}
x = \frac{x_1}{\kappa} + \int_{-\infty}^{x_{-1}} u(x_1, x'_{-1})dx'_{-1} \\
\quad = \frac{x_1}{\kappa} - \ln \frac{g_1}{g_2} , \\
t = x_{-1} .
\end{array}
\right.
$$

This is consistent with the result in [**6, 7**].

Remark 2.11. There are 3 regions in which the above soliton solution becomes regular: (i) $\frac{2}{\kappa} < k_i$, (ii) $-\frac{1}{\kappa} < k_i < \frac{1}{\kappa}$, (iii) $k_i < -\frac{2}{\kappa}$. (Note that this is obtained by the reality condition of e^{ϕ_i}.) In the region (ii), the graph of the soliton solution shows smooth solitons. In the region (i) and (iii), the graph of the soliton solution shows loop solitons.

In Figure 1-4, we show examples of 2-soliton interactions.

3. Conclusions

We have derived the τ-functions of the DP equation by adding the pseudo 3-reduction to the C_∞ 2D-Toda system. We have shown that the τ-functions of the DP equation satisfy the identities of determinants and pfaffians. The result in this article is consistent with the one obtained by Matsuno [**6, 7**]. Our approach clarifies the relationship between the DP equation and the 2D-Toda system.



Figure 1. 2-soliton interaction. $\kappa = 1$, $k_1 = -\frac{1}{3}$, $k_2 = \frac{2}{3}$.



Figure 2. 2-loop soliton interaction. $\kappa = 1$, $k_1 = -\frac{7}{3}$, $k_2 = \frac{8}{3}$.

Appendix A

Let $A = (a_{i,j})_{1 \le i,j \le 2N}$ be a $2N \times 2N$ skew-symmetric matrix, i.e. $a_{i,j} = -a_{j,i}$. The pfaffian of A, $\mathrm{pf}(A)$, is defined as follows:

$$\mathrm{pf}(A) = \mathrm{pf}(a_{i,j})_{1 \le i,j \le 2N} = \left| \begin{array}{cccc} a_{1,2} & a_{1,3} & \cdots & a_{1,2N} \\ & a_{2,3} & \cdots & a_{2,2N} \\ & & \ddots & \vdots \\ & & & a_{2N-1,2N} \end{array} \right. \tag{A.1}$$

$$= \sum_{\sigma} \mathrm{sgn}(\sigma) \prod_{k=1}^{N} a_{i_{2k-1},i_{2k}} \,,$$



FIGURE 3. 2-loop soliton interaction. $\kappa = 1$, $k_1 = \frac{11}{5}$, $k_2 = \frac{10}{3}$.



FIGURE 4. Soliton and loop-soliton interaction. $\kappa = 1$, $k_1 = \frac{7}{3}$, $k_2 = \frac{1}{2}$.

where the summation is taken over all permutations

$$\sigma = \begin{pmatrix} 1 & 2 & \cdots & n \\ i_1 & i_2 & \cdots & i_n \end{pmatrix},$$

satisfying $i_1 < i_2, i_3 < i_4, \cdots, i_{2N-1} < i_{2N}$ and $i_1 < i_3 < \cdots < i_{2N-1}$, and $\mathrm{sgn}(\sigma)$ denotes the parity of the permutation σ.

The pfaffian can be computed recursively by

$$\mathrm{pf}(A) = \sum_{i=2}^{2N} (-1)^i a_{1i} \mathrm{pf}(A_{\hat{1}\hat{i}}), \tag{A.2}$$

where $A_{\hat{1}\hat{i}}$ denotes the matrix A with both the first and i-th rows and columns removed.

The determinant of a skew-symmetric matrix A is the square of the pfaffian of A:

$$\det(A) = [\mathrm{pf}(A)]^2. \tag{A.3}$$

Appendix B

For a bordered determinant, we have the following identity:

$$\begin{vmatrix} \alpha_{1,1} & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 \\ \alpha_{2,1} & \alpha_{2,2} & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ \alpha_{2N,1} & \alpha_{2N,2} & \cdots & \alpha_{2N,2N-1} & \alpha_{2N,2N} & a_{2N} \\ b_1 & b_2 & \cdots & b_{2N-1} & b_{2N} & \delta \end{vmatrix} = \delta^{1-2N} \det |\delta\alpha_{i,j} - a_i b_j|_{1\le i,j\le 2N}, \tag{B.1}$$

where $\delta \neq 0$. This is obtained by adding the $(2N+1)$-th row multiplied by $-a_i/\delta$ to the ith row.

Appendix C

Let $A = (a_{i,j})_{1\le i,j\le 2N+2}$ be a $(2N+2)\times(2N+2)$ skew-symmetric matrix. Assume $a_{2N+1,2N+2} \neq 0$. Adding the $(2N+2)$-th row multiplied by $a_{i,2N+1}/a_{2N+1,2N+2}$ and the $(2N+1)$-th row multiplied by $-a_{i,2N+2}/a_{2N+1,2N+2}$ to the ith row, we obtain

$$\begin{aligned} \det(A) &= \det(a_{i,j})_{1\le i,j\le 2N+2} \\ &= (a_{2N+1,2N+2})^{2-2N} \det(a_{2N+1,2N+2}\, a_{i,j} - a_{2N+1,i}\, a_{2N+2,j} + a_{2N+2,i}\, a_{2N+1,j}), \end{aligned}$$

and

$$\begin{aligned} \mathrm{pf}(A) &= \mathrm{pf}(a_{i,j})_{1\le i,j\le 2N+2} \\ &= (a_{2N+1,2N+2})^{1-N} \mathrm{pf}(a_{2N+1,2N+2}\, a_{i,j} - a_{2N+1,i}\, a_{2N+2,j} + a_{2N+2,i}\, a_{2N+1,j}), \end{aligned}$$

(See e.g. [**22**].) Thus we have the formulae

(C.1)

$$\left|\begin{array}{cccccc} 0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 & b_1 \\ -\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots \\ -\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\ -\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} & b_{2N} \\ -a_1 & -a_2 & \cdots & -a_{2N-1} & -a_{2N} & 0 & \delta \\ -b_1 & -b_2 & \cdots & -b_{2N-1} & -b_{2N} & -\delta & 0 \end{array}\right|$$

$$= \delta^{2-2N} \left|\begin{array}{ccccc} 0 & \beta_{1,2} & \cdots & \beta_{1,2N-1} & \beta_{1,2N} \\ -\beta_{1,2} & 0 & \cdots & \beta_{2,2N-1} & \beta_{2,2N} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -\beta_{1,2N-1} & -\beta_{2,2N-1} & \cdots & 0 & \beta_{2N-1,2N} \\ -\beta_{1,2N} & -\beta_{2,2N} & \cdots & -\beta_{2N-1,2N} & 0 \end{array}\right| ,$$

and

$$\text{(C.2)} \qquad \left|\begin{array}{cccccc} \alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N} & a_1 & b_1 \\ & \alpha_{2,3} & \cdots & \alpha_{2,2N} & a_2 & b_2 \\ & & \ddots & \vdots & \vdots & \vdots \\ & & & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\ & & & & a_{2N} & b_{2N} \\ & & & & & \delta \end{array}\right.$$

$$= \delta^{1-N} \left|\begin{array}{cccc} \beta_{1,2} & \beta_{1,3} & \cdots & \beta_{1,2N} \\ & \beta_{2,2} & \cdots & \beta_{2,2N} \\ & & \ddots & \vdots \\ & & & \beta_{2N-1,2N} \end{array}\right. ,$$

where $\beta_{i,j} = \delta\alpha_{i,j} - a_i b_j + a_j b_i$ and $\delta \neq 0$.

Appendix D

For any determinant A of order n, we have the Jacobi identity

$$A_{ij}A_{pq} - A_{iq}A_{pj} = AA_{ip,jq} , \tag{D.1}$$

where A_{ij} is a first minor which can be obtained by deleting the ith row and the jth column from A, and $A_{ip,jq}$ is a second minor which can be obtained by deleting the ith and pth rows and the jth and qth columns. Setting $r = i = j$ and $s = p = q$, we have

$$A_{rr}A_{ss} - A_{rs}A_{sr} = AA_{rs,rs} . \tag{D.2}$$

Let A be a skew-symmetric determinant. If n is even, then $A_{rr} = A_{ss} = 0$ and $A_{rs} = -A_{sr}$. Therefore we obtain

$$A_{rs}^2 = AA_{rs,rs} , \tag{D.3}$$

which leads to

$$A_{rs} = \mathrm{pf}(A)\mathrm{pf}(A_{rs,rs}) . \tag{D.4}$$

If n is odd, then $A = 0$ and $A_{rs} = A_{sr}$. Therefore we obtain

$$A_{rs}^2 = A_{rr}A_{ss}\,, \tag{D.5}$$

which leads to

$$A_{rs} = \mathrm{pf}(A_{rr})\mathrm{pf}(A_{ss})\,. \tag{D.6}$$

(See [**23, 24**].)

Let

$$A = \begin{vmatrix} 0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 & b_1 \\ -\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \\ -\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\ -\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} & b_{2N} \\ -a_1 & -a_2 & \cdots & -a_{2N-1} & -a_{2N} & 0 & \delta \\ -b_1 & -b_2 & \cdots & -b_{2N-1} & -b_{2N} & -\delta & 0 \end{vmatrix}.$$

Using (D.4) and (C.2), we obtain the formula

(D.7)

$$\begin{vmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & b_1 \\ -\alpha_{1,2} & 0 & \alpha_{2,3} & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & b_2 \\ -\alpha_{1,3} & -\alpha_{2,3} & 0 & \cdots & \alpha_{3,2N-1} & \alpha_{3,2N} & b_3 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots & \vdots \\ -\alpha_{1,2N-1} & -\alpha_{2,2N-1} & -\alpha_{3,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & b_{2N-1} \\ -\alpha_{1,2N} & -\alpha_{2,2N} & -\alpha_{3,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & b_{2N} \\ -a_1 & -a_2 & -a_3 & \cdots & -a_{2N-1} & -a_{2N} & \delta \end{vmatrix}$$

$$= \mathrm{pf}(\alpha_{i,j})_{1\le i,j\le 2N} \begin{vmatrix} \alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N} & a_1 & b_1 \\ & \alpha_{2,3} & \cdots & \alpha_{2,2N} & a_2 & b_2 \\ & & \ddots & \vdots & \vdots & \vdots \\ & & & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\ & & & & a_{2N} & b_{2N} \\ & & & & & \delta \end{vmatrix}.$$

Appendix E

Let

$$A = \begin{vmatrix} 0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 & b_1 \\ -\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 & b_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \\ -\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\ -\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} & b_{2N} \\ c_1 & c_2 & \cdots & c_{2N-1} & c_{2N} & \alpha & \beta \\ d_1 & d_2 & \cdots & d_{2N-1} & d_{2N} & \gamma & \delta \end{vmatrix}.$$

Using the Jacobi identity (D.1), we obtain

$$
\begin{aligned}
&\begin{vmatrix}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 & b_1 \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 & b_2 \\
\vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} & b_{2N} \\
c_1 & c_2 & \cdots & c_{2N-1} & c_{2N} & \alpha & \beta \\
d_1 & d_2 & \cdots & d_{2N-1} & d_{2N} & \gamma & \delta
\end{vmatrix} \\
&\times \begin{vmatrix}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} \\
\vdots & \vdots & \ddots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0
\end{vmatrix} \\
&= \begin{vmatrix}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 \\
\vdots & \vdots & \ddots & \vdots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} \\
c_1 & c_2 & \cdots & c_{2N-1} & c_{2N} & \alpha
\end{vmatrix} \\
&\times \begin{vmatrix}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & b_1 \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & b_2 \\
\vdots & \vdots & \ddots & \vdots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & b_{2N-1} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & b_{2N} \\
d_1 & d_2 & \cdots & d_{2N-1} & d_{2N} & \delta
\end{vmatrix} \\
&- \begin{vmatrix}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & b_1 \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & b_2 \\
\vdots & \vdots & \ddots & \vdots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & b_{2N-1} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & b_{2N} \\
c_1 & c_2 & \cdots & c_{2N-1} & c_{2N} & \beta
\end{vmatrix} \\
&\times \begin{vmatrix}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 \\
\vdots & \vdots & \ddots & \vdots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} \\
d_1 & d_2 & \cdots & d_{2N-1} & d_{2N} & \gamma
\end{vmatrix}.
\end{aligned}
$$

From (D.7), we obtain

$$
\begin{aligned}
&\left|\begin{array}{ccccccc}
0 & \alpha_{1,2} & \cdots & \alpha_{1,2N-1} & \alpha_{1,2N} & a_1 & b_1 \\
-\alpha_{1,2} & 0 & \cdots & \alpha_{2,2N-1} & \alpha_{2,2N} & a_2 & b_2 \\
\vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \\
-\alpha_{1,2N-1} & -\alpha_{2,2N-1} & \cdots & 0 & \alpha_{2N-1,2N} & a_{2N-1} & b_{2N-1} \\
-\alpha_{1,2N} & -\alpha_{2,2N} & \cdots & -\alpha_{2N-1,2N} & 0 & a_{2N} & b_{2N} \\
c_1 & c_2 & \cdots & c_{2N-1} & c_{2N} & \alpha & \beta \\
d_1 & d_2 & \cdots & d_{2N-1} & d_{2N} & \gamma & \delta
\end{array}\right| \\
&= \left|\begin{array}{cccccc}
\alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N} & a_1 & c_1 \\
 & \alpha_{2,3} & \cdots & \alpha_{2,2N} & a_2 & c_2 \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \alpha_{2N-1,2N} & a_{2N-1} & c_{2N-1} \\
 & & & & a_{2N} & c_{2N} \\
 & & & & & \alpha
\end{array}\right| \\
&\times \left|\begin{array}{cccccc}
\alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N} & b_1 & d_1 \\
 & \alpha_{2,3} & \cdots & \alpha_{2,2N} & b_2 & d_2 \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \alpha_{2N-1,2N} & b_{2N-1} & d_{2N-1} \\
 & & & & b_{2N} & d_{2N} \\
 & & & & & \delta
\end{array}\right| \\
&- \left|\begin{array}{cccccc}
\alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N} & b_1 & c_1 \\
 & \alpha_{2,3} & \cdots & \alpha_{2,2N} & b_2 & c_2 \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \alpha_{2N-1,2N} & b_{2N-1} & c_{2N-1} \\
 & & & & b_{2N} & c_{2N} \\
 & & & & & \beta
\end{array}\right| \\
&\times \left|\begin{array}{cccccc}
\alpha_{1,2} & \alpha_{1,3} & \cdots & \alpha_{1,2N} & a_1 & d_1 \\
 & \alpha_{2,3} & \cdots & \alpha_{2,2N} & a_2 & d_2 \\
 & & \ddots & \vdots & \vdots & \vdots \\
 & & & \alpha_{2N-1,2N} & a_{2N-1} & d_{2N-1} \\
 & & & & a_{2N} & d_{2N} \\
 & & & & & \gamma
\end{array}\right| .
\end{aligned}
\tag{E.1}
$$

References

[1] A. Degasperis and M. Procesi, *Asymptotic integrability*, in *Symmetry and Perturbation Theory*, eds. Degasperis and G. Gaeta, World Scientific, Singapore, 1999, pp. 23–37.

[2] A. Degasperis, D. D. Holm and A. N. W. Hone, *A new integrable equation with peakon solutions*, Theoret. and Math. Phys. **133** (2002), 1463–1474.

[3] A. N. W. Hone and J. P. Wang, *Prolongation algebras and Hamiltonian operators for peakon equations*, Inverse Problems **19** (2003), 129–145.

[4] H. Lundmark and J. Szmigielski, *Multi-peakon solutions of the Degasperis-Procesi equation*, Inverse Problems **19** (2003), 1241–1245.

[5] H. Lundmark, *Formation and Dynamics of Shock Waves in the Degasperis-Procesi Equation*, J. Nonl. Sci. **17** (2007), 169–198.

[6] Y. Matsuno, *Multisoliton solution of the Degasperis-Procesi equation and their peakon limit*, Inverse Problems **21** (2005), 1553–1570.

[7] Y. Matsuno, *The N-soliton solution of the Degasperis-Procesi equation*, Inverse Problems **21** (2005), 2085–2101.

[8] A. Constantin and D. Lannes, *The Hydrodynamical Relevance of the Camassa-Holm and Degasperis-Procesi Equations*, Arch. Ration. Mech. Anal. **192** (2009), 165–186.

[9] A. N. Leznov and M. V. Saveliev, *Representation of zero curvature for the system of nonlinear partial differential equations* $x_{a,z,\bar{z}} = \exp(kx)_a$ *and its integrability*, Lett. Math. Phys. **3** (1979), 489–494.

[10] A. V. Mikhailov, *On the integrability of a two-dimensional generalization the Toda lattice*, JETP Lett. **30** (1979), 414–418.

[11] A. V. Mikhailov, *The reduction problem and the inverse scattering method*, Physica **3D** (1981), 73–117.

[12] A. V. Mikhailov *Two-dimensional generalized Toda lattice*, Commun. Math. Phys. **79** (1981), 473–488.

[13] K. Ueno and K. Takasaki, *Toda lattice hierarchy*, Adv. Stud. Pure Math. **4** (1984), 1–95.

[14] V. M. Babich, V. B. Matveev and M. A. Sall, *Binary Darboux transformation for the Toda lattice*, J. Math. Sci. **35** (1986), 2582–2589.

[15] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, 2004.

[16] J. J. C. Nimmo and R. Willox, *Darboux transformations for the two-dimensional Toda system*, Proc. R. Soc. Lond. **453** (1997), 2497–2525.

[17] M. Jimbo and T. Miwa, *Solitons and infinite dimensional Lie algebras*, Publ. RIMS. Kyoto Univ. **19** (1983), 943–1001.

[18] R. Willox, *On a generalized Tzitzeica equation*, Glasgow Math. J. **47A** (2005), 221–231.

[19] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, *KP hierarchies of orthogonal and symplectic type –Transformation groups for soliton equations VI–*, J. Phys. Soc. Japan **50** (1981), 3813–3818.

[20] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, *Transformation groups for soliton equations–Euclidean Lie algebras and reduction of the KP hierarchy*, Publ. RIMS. Kyoto Univ. **18** (1982), 1077–1110.

[21] R. Hirota, *Reduction of soliton equations in bilinear form*, Physica D **18** (1986), 161–170.

[22] I. Satake, *Linear Algebra*, Marcel Dekker Inc., New York, 1975.

[23] T. Muir, *A Treatise on the Theory of Determinants*, Dover, New York, 1960.

[24] R. Vein and P. Dale, *Determinants and Their Applications in Mathematical Physics*, Springer-Verlag, New York, 1999.

Department of Mathematics, The University of Texas - Pan American, Edinburg, Texas 78539-2999

E-mail address: `feg@utpa.edu`

Department of Mathematics, The University of Texas - Pan American, Edinburg, Texas 78539-2999

E-mail address: `kmaruno@utpa.edu`

Department of Mathematics, Kobe University, Rokko, Kobe 657-8501, Japan

E-mail address: `ohta@math.sci.kobe-u.ac.jp`